

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, โรงพยาบาลราชบุรี, มกราคม - ธันวาคม 2022

Organism	TOTAL ISOLATES	BETA - LACTAMS												CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS					
		PENICILLIN	AMPICILLIN	AMOXICILLIN / CLAVULAMIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAZONE	CEFEPIME	OXACILIN	CEFOXITIN	IMPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Typhoidal	1		100 (1)													100 (1)	100 <sup>e</sup> (1)									100 (1)	
<i>Salmonella</i> , Non-typhoidal	116		56.0 (116)													69.0 (116)	68.1 <sup>e</sup> (116)									81.0 (116)	
<i>Shigella</i> spp.			-																								
<i>Staphylococcus aureus</i>		-										- f							-	-	- e	-	-	-	-	-	-
<i>Vibrio cholerae</i> (all serotypes)			-																								
(O1)			-																								
(non O1, non O139)			-																								
<i>Vibrio parahaemolyticus</i>	8		0.0 (8)	100 (8)	100 (8)	100 (8)	- e	25.0 (8)	100 (8)	100 (8)		100 (8)	100 (8)	100 (8)	100 (8)	100 (8)	100 (8)	100 (8)	100 (8)							100 (8)	100 (8)
<i>Vibrio</i> spp.	3		100 (3)	100 (3)	100 (3)	100 (3)	- e	66.7 (3)	100 (3)	100 (3)		100 (3)	66.7 (3)	100 (3)	100 (3)	100 (3)	100 (3)	100 (3)	100 (3)							100 (3)	100 (2)

\* ข้อมูลความไวจากวิธี disk diffusion และ MIC รวมกัน

