

Percentage of susceptible Organisms Isolated From Stool and Rectal Swab, โรงพยาบาลราชบุรี, มกราคม - ธันวาคม 2023

Organism	TOTAL ISOLATES	BETA - LACTAMS												CARBAPENEMS		QUINOLONES		AMINOGLYCOSIDES		GLYCOPEPTIDES		MISCELLANEOUS					
		PENICILLIN	AMPICILLIN	AMOXICILLIN / CLAVULANIC ACID	AMPICILLIN / SULBACTAM	PIPERACILLIN / TAZOBACTAM	CEFAZOLIN (A)	CEFUROXIME SODIUM (parenteral)	CEFOTAXIME	CEFTAZIDIME	CEFTRIAXONE	CEFEPIME	OXACILIN	CEFOXITIN	IMIPENEM	MEROPENEM	CIPROFLOXACIN	LEVOFLOXACIN	AMIKACIN	GENTAMICIN	VANCOMYCIN BY MIC	TEICoplanin	CLINDAMYCIN	ERYTHROMYCIN	CHLORAMPHENICOL	CO-TRIMOXAZOLE	TETRACYCLINE
<i>Salmonella</i> , Typhoidal			-														-	- e								-	
<i>Salmonella</i> , Non-typhoidal	171		52.6 (171)														58.5 (171)	58.5 ^e (171)								82.4 (170)	
<i>Shigella</i> spp.			-														-	-								-	
<i>Staphylococcus aureus</i>		-										- f					-	-	-	- e	- e	-	-	-	-	-	-
<i>Vibrio cholerae</i> (all serotypes)	1		100 (1)																						-	100 (1)	-
(O1)	1		100 (1)																						-	100 (1)	-
(non O1, non O139)			-																						-	-	-
<i>Vibrio parahaemolyticus</i>	12		0.0 (12)	100 (12)	100 (12)	100 (12)	- e	50.0 (12)	100 (12)	100 (12)		100 (12)	100 (12)	100 (12)	100 (12)	100 (12)	100 (12)	100 (12)	100 (12)							100 (12)	-
<i>Vibrio</i> spp.	5		60.0 (5)	80.0 (5)	60.0 (5)	100 (5)	- e	80.0 (5)	100 (5)	100 (5)		100 (5)	40.0 (5)	100 (5)	80.0 (5)	60.0 (5)	60.0 (5)	100 (5)	80.0 (5)							60.0 (5)	-

* ข้อมูลความไวจากวิธี disk diffusion และ MIC รวมกัน

